


Celiac Disease

What is celiac disease?

Celiac disease is a digestive disease that damages the small intestine and interferes with absorption of nutrients from food. People who have celiac disease cannot tolerate a protein called gluten, found in wheat, rye, and barley. Gluten is found mainly in foods but may also be found in products we use every day, such as stamp and envelope adhesive, medicines, and vitamins.

When people with celiac disease eat foods or use products containing gluten, their immune system responds by damaging the small intestine. The tiny, fingerlike protrusions lining the small intestine are damaged or destroyed. Called villi, they normally allow nutrients from food to be absorbed into the bloodstream. Without healthy villi, a person becomes malnourished, regardless of the quantity of food eaten.



Because the body's own immune system causes the damage, celiac disease is considered an autoimmune disorder. However, it is also classified as a disease of malabsorption because nutrients are not absorbed. Celiac disease is also known as celiac sprue, nontropical sprue, and gluten-sensitive enteropathy. Celiac disease is a genetic disease, meaning it runs in families. Sometimes the disease is triggered—or becomes active for the first time—after surgery, pregnancy, childbirth, viral infection, or severe emotional stress.

What are the symptoms of celiac disease?

Celiac disease affects people differently. Symptoms may occur in the digestive system, or in other parts of the body. For example, one person might have diarrhea and abdominal pain, while another person may be irritable or depressed. In fact, irritability is one of the most common symptoms in children.

Symptoms of celiac disease may include one or more of the following:

- gas
- recurring abdominal bloating and pain
- chronic diarrhea
- constipation
- pale, foul-smelling, or fatty stool
- weight loss/weight gain
- fatigue
- unexplained anemia (a low count of red blood cells causing fatigue)
- bone or joint pain
- osteoporosis, osteopenia
- behavioral changes
- tingling numbness in the legs (from nerve damage)
- muscle cramps
- seizures
- missed menstrual periods (often because of excessive weight loss)
- infertility, recurrent miscarriage
- delayed growth
- failure to thrive in infants
- pale sores inside the mouth, called aphthous ulcers
- tooth discoloration or loss of enamel
- itchy skin rash called dermatitis herpetiformis

A person with celiac disease may have no symptoms. People without symptoms are still at risk for the



Celiac Disease

complications of celiac disease, including malnutrition. The longer a person goes undiagnosed and untreated, the greater the chance of developing malnutrition and other complications. Anemia, delayed growth, and weight loss are signs of malnutrition: The body is just not getting enough nutrients. Malnutrition is a serious problem for children because they need adequate nutrition to develop properly. (See [Complications](#).)

How is celiac disease diagnosed?

Recognizing celiac disease can be difficult because some of its symptoms are similar to those of other diseases. In fact, sometimes celiac disease is confused with irritable bowel syndrome, iron-deficiency anemia caused by menstrual blood loss, Crohn's disease, diverticulitis, intestinal infections, and chronic fatigue syndrome. As a result, celiac disease is commonly underdiagnosed or misdiagnosed.

Recently, researchers discovered that people with celiac disease have higher than normal levels of certain autoantibodies in their blood. Antibodies are protective proteins produced by the immune system in response to substances that the body perceives to be threatening. Autoantibodies are proteins that react against the body's own molecules or tissues. To diagnose celiac disease, physicians will usually test blood to measure levels of

- Immunoglobulin A (IgA)
- anti-tissue transglutaminase (tTGA)
- IgA anti-endomysium antibodies (AEA)

Before being tested, one should continue to eat a regular diet that includes foods with gluten, such as breads and pastas. If a person stops eating foods with gluten before being tested, the results may be negative for celiac disease even if celiac disease is actually present.

If the tests and symptoms suggest celiac disease, the doctor will perform a small bowel biopsy. During the biopsy, the doctor removes a tiny piece of tissue from the small intestine to check for damage to the villi. To obtain the tissue sample, the doctor eases a long, thin tube called an endoscope through the mouth and stomach into the small intestine. Using instruments passed through the endoscope, the doctor then takes the sample.

What is the treatment?

The only treatment for celiac disease is to follow a gluten-free diet. When a person is first diagnosed with celiac disease, the doctor usually will ask the person to work with a dietitian on a gluten-free diet plan. A dietitian is a health care professional who specializes in food and nutrition. Someone with celiac disease can learn from a dietitian how to read ingredient lists and identify foods that contain gluten in order to make informed decisions at the grocery store and when eating out.

For most people, following this diet will stop symptoms, heal existing intestinal damage, and prevent further damage. Improvements begin within days of starting the diet. The small intestine is usually completely healed in 3 to 6 months in children and younger adults and within 2 years for older adults. Healed means a person now has villi that can absorb nutrients from food into the bloodstream.

In order to stay well, people with celiac disease must avoid gluten for the rest of their lives. Eating any gluten, no matter how small an amount, can damage the small intestine. The damage will occur in anyone with the disease, including people without noticeable symptoms. Depending on a person's age at diagnosis, some problems will not improve, such as delayed growth and tooth discoloration.

Some people with celiac disease show no improvement on the gluten-free diet. This condition is called unresponsive celiac disease. The most common reason for poor response is that small amounts of gluten are still present in the diet. Advice from a dietitian who is skilled in educating patients about the gluten-free diet is essential to achieve the best results.

Rarely, the intestinal injury will continue despite a strictly gluten-free diet. People in this situation have severely damaged intestines that cannot heal. Because their intestines are not absorbing enough nutrients, they may need to receive nutrients directly into their bloodstream through a vein, or intravenously. People with this condition may need to be evaluated for complications of the disease. Researchers are



Celiac Disease

now evaluating drug treatments for unresponsive celiac disease.

The web contains information about celiac disease, some of which is not accurate. The best people for advice about diagnosing and treating celiac disease are one's doctor and dietitian.

The Gluten-free Diet

A gluten-free diet means not eating foods that contain wheat (including spelt, triticale, and kamut), rye, and barley. The foods and products made from these grains are also not allowed. In other words, a person with celiac disease should not eat most grain, pasta, cereal, and many processed foods. Despite these restrictions, people with celiac disease can eat a well-balanced diet with a variety of foods, including gluten-free bread and pasta. For example, people with celiac disease can use potato, rice, soy, amaranth, quinoa, buckwheat, or bean flour instead of wheat flour. They can buy gluten-free bread, pasta, and other products from stores that carry organic foods, or order products from special food companies. Gluten-free products are increasingly available from regular stores.

Checking labels for "gluten free" is important since many corn and rice products are produced in factories that also manufacture wheat products. Hidden sources of gluten include additives such as modified food starch, preservatives, and stabilizers. Wheat and wheat products are often used as thickeners, stabilizers, and texture enhancers in foods.

"Plain" meat, fish, rice, fruits, and vegetables do not contain gluten, so people with celiac disease can eat as much of these foods as they like. Recommending that people with celiac disease avoid oats is controversial because some people have been able to eat oats without having symptoms. Scientists are currently studying whether people with celiac disease can tolerate oats. Until the studies are complete, people with celiac disease should follow their physician's or dietitian's advice about eating oats. Examples of foods that are safe to eat and those that are not are provided in the [table](#) below.

The gluten-free diet is challenging. It requires a completely new approach to eating that affects a person's entire life. Newly diagnosed people and their families may find support groups to be particularly helpful as they learn to adjust to a new way of life. People with celiac disease have to be extremely careful about what they buy for lunch at school or work, what they purchase at the grocery store, what they eat at restaurants or parties, or what they grab for a snack. Eating out can be a challenge. If a person with celiac disease is in doubt about a menu item, ask the waiter or chef about ingredients and preparation, or if a gluten-free menu is available.

Gluten is also used in some medications. One should check with the pharmacist to learn whether medications used contain gluten. Since gluten is also sometimes used as an additive in unexpected products, it is important to read all labels. If the ingredients are not listed on the product label, the manufacturer of the product should provide the list upon request. With practice, screening for gluten becomes second nature.

The Gluten-free Diet: Some Examples

In 2006, the American Dietetic Association updated its recommendations for a gluten-free diet. The following chart is based on the 2006 recommendations. This list is **not** complete, so people with celiac disease should discuss gluten-free food choices with a dietitian or physician who specializes in celiac disease. People with celiac disease should always read food ingredient lists carefully to make sure that the food does not contain gluten.



Celiac Disease

Allowed Foods		
Amaranth	Job's tears	Sago
Arrowroot	Legumes	Seeds
Buckwheat	Millet	Soy
Cassava	Nuts	Sorghum
Corn	Potatoes	Tapioca
Flax	Quinoa	Wild Rice
Indian rice grass	Rice	Yucca
Foods To Avoid		
Wheat	Barley	
<ul style="list-style-type: none"> Including einkorn, emmer, spelt, kamut Wheat starch, wheat bran, wheat germ, cracked wheat, hydrolyzed wheat protein 	Rye	
	Triticale (a cross between wheat and rye)	
Other Wheat Products		
Bromated flour	Graham flour	Self-rising flour
Durum flour	Phosphated flour	Semolina
Enriched flour	Plain flour	White flour
Farina		
Processed Foods That May Contain Wheat, Barley, or Rye*		
Bouillon cubes	French fries	Seasoned tortilla chips
Brown rice syrup	Gravy	Self-basting turkey
Chips/potato chips	Imitation fish	Soups
Candy	Matzo	Soy sauce
Cold cuts, hot dogs, salami, sausage	Rice mixes	Vegetables in sauce
Communion wafer	Sauces	

* Most of these foods can be found gluten-free. When in doubt, check with the food manufacturer.



Celiac Disease

From the following resource: Thompson T. *Celiac Disease Nutrition Guide*, 2nd ed. Chicago: American Dietetic Association; 2006. © American Dietetic Association. Adapted with permission. For a complete copy of the *Celiac Disease Nutrition Guide*, please visit www.eatright.org.

What are the complications of celiac disease?

Damage to the small intestine and the resulting nutrient absorption problems put a person with celiac disease at risk for malnutrition, anemia, and several other diseases and health problems.

- **Lymphoma and adenocarcinoma** are cancers that can develop in the intestine.
- **Osteoporosis** is a condition in which the bones become weak, brittle, and prone to breaking. Poor calcium absorption contributes to osteoporosis.
- **Miscarriage and congenital malformation** of the baby, such as neural tube defects, are risks for pregnant women with untreated celiac disease because of nutrient absorption problems.
- **Short stature** refers to being significantly under the average height. Short stature results when childhood celiac disease prevents nutrient absorption during the years when nutrition is critical to a child's normal growth and development. Children who are diagnosed and treated before their growth stops may have a catch-up period.

How common is celiac disease?

Data on the prevalence of celiac disease is spotty. In Italy about 1 in 250 people, and in Ireland about 1 in 300 people, have celiac disease. Recent studies have shown that it may be more common in Africa, South America, and Asia than previously believed.

Until recently, celiac disease was thought to be uncommon in the United States. However, studies have shown that celiac disease is very common. Recent findings estimate about 2 million people in the United States have celiac disease, or about 1 in 133 people. Among people who have a first-degree relative diagnosed with celiac disease, as many as 1 in 22 people may have the disease.

Celiac disease could be underdiagnosed in the United States for a number of reasons including:

- Celiac symptoms can be attributed to other problems.
- Many doctors and health care providers are not knowledgeable about the disease.
- Only a small number of U.S. laboratories are experienced and skilled in testing for celiac disease.

More research is needed to learn the true prevalence of celiac disease among Americans.

Diseases Linked to Celiac Disease

People with celiac disease tend to have other autoimmune diseases. The connection between celiac disease and these diseases may be genetic. These diseases include

- thyroid disease
- systemic lupus erythematosus
- type 1 diabetes
- liver disease
- collagen vascular disease
- rheumatoid arthritis



Celiac Disease

Sjögren's syndrome

Dermatitis Herpetiformis

Dermatitis herpetiformis (DH) is a severe, itchy, blistering skin manifestation of celiac disease. Not all people with celiac disease develop dermatitis herpetiformis. The rash usually occurs on the elbows, knees, and buttocks. Unlike other forms of celiac disease, the range of intestinal abnormalities in DH is highly variable, from minimal to severe. Only about 20 percent of people with DH have intestinal symptoms of celiac disease.

To diagnose DH, the doctor will test the person's blood for autoantibodies related to celiac disease and will biopsy the person's skin. If the antibody tests are positive and the skin biopsy has the typical findings of DH, patients do not need to have an intestinal biopsy. Both the skin disease and the intestinal disease respond to a gluten-free diet and recur if gluten is added back into the diet. In addition, the rash symptoms can be controlled with medications such as dapsone (4',4'diamino-diphenylsufone). However, dapsone does not treat the intestinal condition and people with DH should also maintain a gluten-free diet.

Points to Remember

- People with celiac disease cannot tolerate gluten, a protein in wheat, rye, barley, and possibly oats.
- Untreated celiac disease damages the small intestine and interferes with nutrient absorption.
- A person with celiac disease may or may not have symptoms.
- Since celiac disease is hereditary, family members of a person with celiac disease may wish to be tested.
- Celiac disease is treated by eliminating all gluten from the diet. The gluten-free diet is a lifetime requirement.
- A dietitian can teach a person with celiac disease food selection, label reading, and other strategies to help manage the disease.

The U.S. Government does not endorse or favor any specific commercial product or company. Trade, proprietary, or company names appearing in this document are used only because they are considered necessary in the context of the information provided. If a product is not mentioned, the omission does not mean or imply that the product is unsatisfactory.

For More Information

American Dietetic Association

120 South Riverside Plaza, Suite 2000
Chicago, IL 60606-6995
Phone: 1-800-877-1600 or 312-899-0040
Fax: 312-899-4739
Email: hotline@eatright.org
Internet: www.eatright.org

Celiac Disease Foundation

13251 Ventura Boulevard, #1
Studio City, CA 91604
Phone: 818-990-2354
Fax: 818-990-2379
Email: cdf@celiac.org
Internet: www.celiac.org



Celiac Disease

Celiac Sprue Association/USA Inc.

P.O. Box 31700

Omaha, NE 68131-0700

Phone: 1-877-CSA-4CSA (272-4272)

Fax: 402-558-1347

Email: celiacs@csaceliacs.org

Internet: www.csaceliacs.org